Ratio between two Λ and $\bar{\Lambda}$ production mechanisms in p scattering

B. Hoeneisen[1]

[1] *Universidad San Francisco de Quito, Quito, Ecuador*

(Dated: April 18, 2016)

We consider Λ and $\bar{\Lambda}$ production in a wide range of proton scattering experiments. The produced Λ and $\bar{\Lambda}$ may or may not contain a diquark remnant of the beam proton. The ratio of these two production mechanisms is found to be a simple universal function $r = [\kappa/(y_p - y)]^i$ of the rapidity difference $y_p - y$ of the beam proton and the produced Λ or $\bar{\Lambda}$, valid over four orders of magnitude, from $r \approx 0.01$ to $r \approx 100$, with $\kappa = 2.86 \pm 0.03 \pm 0.07$, and $i = 4.39 \pm 0.06 \pm 0.15$.

arXiv:1604.05379v1 [hep-ph] 18 Apr 2016

The $\bar{\Lambda}/\Lambda$ production ratio measured in a wide range of proton scattering experiments $pZ \to \Lambda(\bar{\Lambda})X$ has been found to be a universal function $f(y_p - y)$ of "rapidity loss" $y_p - y$, where y_p and y are, respectively, the rapidities of the beam proton and the produced Λ or $\bar{\Lambda}$ [1, 2]. The function $f(y_p - y)$ is observed to be independent, or depends only weakly, on the total center of mass energy $\sqrt{s}$ of the two colliding hadrons in the range 0.024 to 7 TeV, on the target $Z = p, \bar{p}$, Be or Pb, on the transverse momentum p_T of the Λ or $\bar{\Lambda}$, or on sample composition [2]. We consider the picture in which an s quark produced in the scattering may coalesce with a ud diquark remnant of the beam proton and produce a Λ [2–6].

Let $n_\Lambda(y)$ and $n_{\bar{\Lambda}}(y)$ be the distributions of Λ's and $\bar{\Lambda}$'s as a function of rapidity y in the center of mass frame of the two colliding hadrons. Rapidity is defined so that the p beam has positive rapidity y_p. We write these distributions as follows:

$$\begin{aligned} n_\Lambda(y) &= n_{\Lambda\alpha1}(y) + n_{\Lambda\alpha2}(y) + n_{\Lambda\beta}(y), \\ n_{\bar{\Lambda}}(y) &= n_{\bar{\Lambda}\alpha1}(y) + n_{\bar{\Lambda}\alpha2}(y) + n_{\bar{\Lambda}\beta}(y), \end{aligned} \tag{1}$$

where $n_{\Lambda\alpha1}(y)$ is the distribution of Λ's containing a diquark remnant of beam 1, $n_{\Lambda\alpha2}(y)$ is the distribution of Λ's containing a diquark remnant of beam 2, and $n_{\Lambda\beta}(y)$ is the distribution of Λ's containing no beam remnant, and similarly for $\bar{\Lambda}$'s. Production mechanism β has no memory of the beams and hence $n_{\Lambda\beta}(y) = n_{\bar{\Lambda}\beta}(y) \equiv n_\beta(y) = n_\beta(-y)$. The distribution $n_\beta(y)$ is approximately independent of y within the "rapidity plateau" $-y_{\max} < y < y_{\max}$ as shown schematically in Fig. 1. See also Fig. 2 of [2], and Fig. 50.4 of [7]. We assume that beam 1 is a proton beam, so $n_{\bar{\Lambda}\alpha1} = 0$. If beam 2 is a p beam, $n_{\bar{\Lambda}\alpha2} = 0$. If beam 2 is a $\bar{p}$ beam, $n_{\Lambda\alpha2} = 0$.

We now consider events with $y > y_{\min}$ so that $n_{\bar{\Lambda}\alpha2}$ and $n_{\Lambda\alpha2}$ can be neglected. The $\bar{\Lambda}/\Lambda$ production ratio can then be written as:

$$f = \frac{n_\beta(y)}{n_\beta(y) + n_{\Lambda\alpha1}(y)} \equiv \frac{1}{1+r}, \tag{2}$$

where the ratio, r, of the yields of the two production mechanisms α and β is

$$r \equiv \frac{n_{\Lambda\alpha1}(y)}{n_\beta(y)} = \frac{1}{f} - 1. \tag{3}$$



FIG. 1: Distributions of Λ and $\bar{\Lambda}$ production as a function of rapidity y in the center of mass frame of $p\bar{p}$ scattering. The beam rapidities are respectively y_p and $y_{\bar{p}}$. The sub-index α (β) denotes Λ's or $\bar{\Lambda}$'s containing (not containing) a beam diquark remnant. Lowering the total center of mass energy $\sqrt{s}$ translates y_p and $n_{\Lambda\alpha1}(y)$ left, and translates $y_{\bar{p}}$ and $n_{\bar{\Lambda}\alpha2}(y)$ right.

The purpose of this note is to point out that the ratio r can be fit over four orders of magnitude, from $r \approx 0.01$ to $r \approx 100$, with a simple universal function with only two parameters κ and i:

$$r = \left[\frac{\kappa}{y_p - y}\right]^i. \tag{4}$$

Figure 2 presents the ratios $r = 1/f - 1$ measured in a wide range of proton scattering experiments. The data points with $y < 0.75$ of the DØ $p\bar{p}$ experiment were omitted because for them we can not neglect $n_{\bar{\Lambda}\alpha2}(y)$. The data point of the STAR pp experiment has $y = 0$, so $n_{\Lambda\alpha1} = n_{\Lambda\alpha2}$. Therefore we have divided $1/f - 1$ by 2.

The parameters κ and i have a simple interpretation and can be read off the $\log(y_p - y)$ vs $\log r$ graph in Fig. 2: $\kappa \approx 2.8$ is the rapidity loss at which $r = 1$, and $-i \approx -4.4$ is the slope of the straight line in Fig. 2.


ALICE, pp, √s=7 TeV
ALICE, pp, √s=2.76 TeV
ALICE, pp, √s=0.9 TeV
ATLAS, pp, √s=7 TeV
ATLAS, pp, √s=0.9 TeV
DO, pp̄, √s=1.96 TeV
STAR, pp, √s=0.2 TeV
LHCb, pp, √s=7 TeV
LHCb, pp, √s=0.9 TeV
R-607, pp, √s=0.0623 TeV
R-603, pp, √s=0.053 TeV
E8, p Be, √s=0.024 TeV
E8, p Pb, √s=0.024 TeV
r = n_Λα1/n_β production ratio
Rapidity loss Δy = y_p - y

FIG. 2: Ratio $r \equiv n_{\Lambda\alpha 1}(y)/n_\beta(y)$ of Λ's produced with and without a beam proton remnant, as a function of rapidity loss $y_p - y$, obtained by the experiments listed [2, 8–13]. The fit shown has the form $r = [\kappa/(y_p - y)]^i$ with $\kappa = 2.862$ and $i = 4.393$.

For $y_p - y < \kappa$ production mechanism α dominates. For $y_p - y > \kappa$ production mechanism β dominates.

The fit to all of the data in Fig. 2 obtains $\kappa = 2.79 \pm 0.03$ and $i = 4.54 \pm 0.08$ with $\chi^2 = 637$ for 121 degrees of freedom. The large χ^2 is due to tension between the data points of different experiments as can be seen in Fig. 2. Omitting the R-603 and R-607 measurements, which have some data points off the rapidity plateau, obtains $\kappa = 2.86 \pm 0.03$ and $i = 4.39 \pm 0.06$ with $\chi^2 = 342$ for 102 degrees of freedom. This is the fit shown in Fig. 2. Fitting only the E8 Pb data points where production mechanism α dominates obtains $\kappa = 2.93 \pm 0.15$ and $i = 4.06 \pm 0.30$ with $\chi^2 = 10.6$ for 13 degrees of freedom. Fitting all data with $y_p - y > 2.8$, where production mechanism β dominates, obtains $\kappa = 2.94 \pm 0.10$ and $i = 4.23 \pm 0.25$ with $\chi^2 = 37$ for 32 degrees of freedom. In conclusion, we see no significant departure from Eq. (4) at either end of the data range. Our final estimate from several fits is

$$\kappa = 2.86 \pm 0.03 \pm 0.07, i = 4.39 \pm 0.06 \pm 0.15, \qquad (5)$$

where the first uncertainty is statistical from the fit, and the second uncertainty is systematic and accounts for different data selections for the fits.

From $\sqrt{s} = 0.024$ to 7 TeV the cross-sections $\sigma(pp)$ and $\sigma(p\bar{p})$, and the width of the rapidity plateau $2y_{\rm max}$, increase by approximately a factor 2 [7], so $n_\beta(y)$ is approximately independent of $\sqrt{s}$ and y on the rapidity plateau. We conclude that the probability density that a p scatters and becomes a Λ with rapidity y is proportional to $[\kappa/(y_p - y)]^i$. This result should also be valid for Λ_c, Λ_b, Σ^+, etc.

I thank my colleagues in the D0 Collaboration for their comments and inspiration to undertake this analysis.

[1] R. Aaij *et al.* (LHCb Collaboration), Measurement of V^0 production ratios in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV, J. High Energy Phys. 1108 (2011) 034.
[2] V.M. Abazov *et al.* (D0 Collaboration), Measurement of the forward-backward asymmetry of Λ and $\bar{\Lambda}$ production in $p\bar{p}$ collisions, Phys. Rev. D **93**, 032002 (2016).
[3] K. P. Das and Rudolph C. Hwa, Quark-antiquark recombination in the fragmentation region, Phys. Lett. B **68**, 5 (1977).
[4] Stanley J. Brodsky and John F. Gunion, Hadronic fragmentation as a probe of the underlying dynamics of hadron collisions, Phys. Rev. D **17**, 3 (1978).
[5] D. Cutts *et al.*, Experimental study of low-p_t hadron fragmentation, Phys. Rev. Lett. 43, 5 (1979).
[6] Rudolph C. Hwa and R. G. Roberts, Pion structure function from low-p_t hadron production, Z. Physik C, **1**, 81-86 (1979) and references therein.
[7] K.A. Olive *et al.* (Particle Data Group), Review of particle physics, Chin. Phys. C, 36, 090001 (2014)
[8] E. Abbas *et al.* (ALICE Collaboration), Mid-rapidity anti-baryon to baryon ratios in pp collisions at $\sqrt{s} = 0.9, 2.76,$ and 7 TeV measured by ALICE, Eur. Phys J. C 73, 2496 (2013).
[9] G. Aad *et al.* (ATLAS Collaboration), K^0_S and Λ production in pp interactions at $\sqrt{s} = 0.9$ and 7 TeV measured with the ATLAS detector at the LHC, Phys. Rev. D **85** (2012) 012001.
[10] B.I. Abelev *et al.* (STAR Collaboration), Strange particle production in $p + p$ collisions at $\sqrt{s} = 200$ GeV, Phys. Rev. C **75**, 064901 (2007).
[11] G.J. Bobbink *et al.* (R-607 Collaboration), The production of high-momentum particles and resonances in pp collisions at the CERN intersecting storage rings, Nucl. Phys. B **217** (1983) 11.
[12] S. Erhan *et al.* (R-603 Collaboration), Hyperon production in pp interactions at $\sqrt{s} = 53$ and 62 GeV, Phys. Lett. B **85** (1979) 447.
[13] P. Skubic *et al.* (E8 Collaboration), Neutral-strange-particle production by 300-GeV protons, Phys. Rev. D **18**, 3115 (1978).